

## Correlation between immunization status and pediatric diphtheria patients outcomes in the Sampang District, 2011-2015

Kevin Sastra Dhinata<sup>1</sup>, Atika<sup>2</sup>, Dominicus Husada<sup>3</sup>, Dwiyanti Puspitasari<sup>3</sup>

### Abstract

**Background** The number of diphtheria cases recently increased, such that an outbreak was declared in East Java Province, which includes the Sampang District. Immunization completion status is a determining factor for diphtheria infection.

**Objective** To investigate for correlations between immunization status and outcomes (severity level, fatality, and complications) of diphtheria patients in the Sampang District.

**Methods** This analytic, cross-sectional study used secondary data from the East Java Provincial Health Office on diphtheria patients aged 0-20 years during the 2011-2015 outbreak in the Sampang District and interviews with diphtheria patients in that region. The Diphtheria Research Team of Dr. Soetomo Hospital collected data on immunization status, diphtheria severity (mild, moderate, or severe), case fatality (died or survived), and complications in the patients (with or without complications). Spearman's, Chi-square, and Fisher's exact tests were used for data analyses, accordingly.

**Results** Seventy-one patients with clinical diphtheria were identified, 17 of whom were confirmed with positive culture results. The case fatality rates were 7% in patients with clinical and 5.9% in confirmed diphtheria. There were no correlations between patient immunization status and severity ( $P=0.469$  clinical,  $P=0.610$  confirmed), or fatality ( $P=0.618$  clinical,  $P=0.294$  confirmed) of diphtheria in the clinical and confirmed diphtheria patients. However, there was a correlation between patient immunization status and the emergence of complications in clinical ( $P=0.013$ ), but not in confirmed ( $P=0.620$ ) diphtheria patients.

**Conclusion** There is a correlation between immunization status and complications in clinical diphtheria patients. Such a correlation is not found in confirmed diphtheria cases because none of the patients had complete immunization status. [Paediatr Indones. 2018;58:110-15; doi: <http://dx.doi.org/10.14238/pi58.3.2018.110-15>].

**Keywords:** diphtheria; immunization status; diphtheria severity; fatality; complication

Diphtheria is an acute infectious disease that mainly affects infants and children in the early years of life.<sup>1</sup> The number of diphtheria cases has decreased throughout the world. However, the East Java Provincial Government of Indonesia declared a diphtheria outbreak due to a high number of diphtheria cases in the province, including the Sampang District. From 2009 to 2012, there were approximately 1,870 reported diphtheria cases in the province, with 643 cases clinically diagnosed with diphtheria in healthcare centers. Out of 643 cases of diphtheria based on clinical diagnosis, fifty of them were confirmed by positive culture results based on official reports released between 2013 to 2014.<sup>2,3</sup> In the Sampang District alone, approximately 91 cases were found from 2011 to 2015.

Diphtheria is caused by the *Corynebacterium diphtheriae* bacterial exotoxin.<sup>1</sup> The diphtheria toxin's main mode of action is to inhibit protein synthesis. This toxin may spread relatively quickly to different parts of

---

From the Medical School<sup>1</sup>, Department of Public Health and Preventive Medicine<sup>2</sup>, and Department of Child Health<sup>3</sup>, Airlangga University/Dr. Soetomo General Hospital, Surabaya, East Java, Indonesia.

**Reprint requests to:** Kevin Sastra Dhinata. Department of Child Health, Airlangga University Faculty of Medicine/Dr. Soetomo General Hospital, Surabaya, Indonesia.Jl. Prof. Dr. Moestopo No. 6-8, Surabaya, East Java, Indonesia.Email: kevsastr@gmail.com.

the body, causing complications in the patients, such as myocarditis and neurologic dysfunction.<sup>4-6</sup> Diphtheria is transmitted mainly by droplets from sneezing and coughing, but also may be transmitted through direct contact with cutaneous diphtheria wounds.<sup>7</sup> Initially, the signs and symptoms of diphtheria are non-specific, such as malaise, fever around 38° Celsius, hoarseness, sore throat, and rhinorrhea. As the disease progresses, pseudomembranes develop in most cases, which may cause respiratory tract obstruction. This obstruction may lead to patient fatality.<sup>7</sup> A definitive diagnosis of diphtheria is carried out by obtaining specimens from suspected lesions and culturing them in media specific for *Corynebacterium diphtheriae*.<sup>4,6,7</sup> Supportive examinations may also be carried out to assist the diagnostic process, in form of polymerase chain reaction (PCR), antibiotic-sensitivity test to identify antimicrobial resistance, Elek test to identify diphtheria toxin, and Shick test to identify serum antibodies against the diphtheria toxin.<sup>8,9</sup>

The number of diphtheria cases has been significantly reduced since the discovery and introduction of a diphtheria vaccination. Vaccine administration stimulates the production of antibodies against diphtheria toxin, mainly IgG and IgA, making the toxin unable to bind to the toxin receptor during future infection.<sup>4</sup> Studies have shown that of many factors affecting diphtheria spread in society, immunizations stand out as the most significant and influential factor.<sup>10</sup> We aimed to identify correlations between immunization status and diphtheria severity level, fatality, and complications in the Sampang District during the diphtheria outbreak from 2011 to 2015.

## Methods

This study was an analytic study with a cross-sectional approach. Subjects were patients with clinical diagnoses of diphtheria in the Sampang District of Madura Island, East Java. Data were obtained from the East Java Provincial Health Office on diphtheria patients from 2011-2015 and by interview conducted towards the patient of the 2011-2015 diphtheria outbreak in the region by the Diphtheria Research Team of Soetomo Hospital consisting of pediatrician, residents, doctors, epidemiologists, public health

officers, nurses, and medical students. The inclusion criteria were patients with a clinical diagnosis of diphtheria, maximum age of 20 years at the time of diagnosis, complete data and/or medical records, and diagnosed as not immunocompromised by a medical doctor. The minimum required number of subjects was calculated using the hypothesis test formula for two samples, and found to be 56.

The independent variable in this study was the patient immunization status, which was classified as "complete," "incomplete," or "unimmunized," according to the Indonesian Ministry of Health standards.<sup>11</sup> The dependent variables were the degree of severity, fatality, and complications in the patients. The degree of diphtheria severity was classified according to the 2008 *Diagnosis and Therapy Guidelines* of Dr. Soetomo General Hospital, Surabaya, Indonesia, into "severe," "moderate," or "mild" diphtheria. Fatality was defined as the final outcome of death due to diphtheria. Complication was defined as the presence of bullneck, myocarditis, shock, or bleeding manifestation.<sup>6</sup> Spearman's, Chi-square, and Fisher's exact tests were used for statistical analyses by SPSS Statistics 23.0 software. This study was approved by the Ethics Committee for Health Research of Airlangga University Medical School, Surabaya.

## Results

There were 81 diphtheria patients recorded in the Sampang District during 2011-2015. Ten patients were not included because of aged  $\geq 20$  years and/or incomplete data. Seventy-one patients with clinical diagnoses of diphtheria were included in this study, 17 (23.9%) of whom had positive culture results for *Corynebacterium diphtheriae*. Most of them were positive for *mitis biovar*, as described in Table 1.

Most diphtheria patients were female and characteristics of the patients were relatively similar between the two groups (clinical and confirmed

**Table 1.** Culture results of the subjects

Culture results	(N=71)
Positive, n (%)	
Gravis biovar	2 (2.8)
Mitis biovar	15 (21.1)
Negative, n(%)	54 (76.1)

diphtheria). Both groups were predominantly aged 5-9 years at the time of infection. Both groups also had relatively similar age characteristics. But the patients with confirmed diphtheria had a slightly narrower age range at the time of infection. Subjects' characteristics are summarized in **Table 2**.


**Table 2.** Characteristics of the subjects

Characteristics	Clinical diphtheria (n=71)	Confirmed diphtheria (N=17)
Sex, n (%)		
Male	29 (59.2)	7 (41.2)
Female	42 (40.8)	10 (58.8)
Age group, n (%)		
0-4 years	21 (29.6)	4 (23.5)
5-9 years	31 (43.7)	8 (47.1)
10-19 years	19 (26.8)	5 (29.4)
Age characteristics, n (%)		
Mean (SD), years	7.19 (4.289)	7.77 (4.2061)
Median, years	6	7
Minimum, years	1	3
Maximum, years	19	18

Diphtheria patients were found in 12 out of the total of 14 districts in the Sampang District. A large number of cases were found in Banyuates District and Sampang District. However, no patients were found in Sresek and Pangarengan Districts, as shown in **Figure 1**.

There was a larger proportion of severe diphtheria in patients without complete immunizations. However, Spearman's correlation test revealed no significant correlation between immunization status and the degree of diphtheria severity of in patients with clinical and confirmed diphtheria (**Table 3**).

Fisher's exact test revealed no significant



**Figure 1.** Map of diphtheria patient distribution by-district in the Sampang District. Left box (black font): number of patients with clinical diphtheria; Right box (red font): number of patients with confirmed diphtheria.

correlation between immunization status and fatality status in diphtheria patients with clinical and confirmed diphtheria (**Table 4**).

Complications occurred in 26 out of 71 (36.6%) diphtheria patients. The majority of patients with complications had bullneck, as displayed in **Table 5**. From the total of 26 patients having complications of diphtheria, 8 of them belong to the group of 17 patients with confirmed diphtheria, which were analyzed further. Chi-square test revealed a significant correlation between immunization status and complications of patients with clinical diphtheria, but no correlation between immunization status and complications in patients with confirmed diphtheria (**Table 6**).

**Table 3.** Analysis of immunization status and degree of severity

Diphtheria status	Immunization status	Diphtheria severity			P value
		Mild	Moderate	Severe	
Clinical diphtheria	Complete, n(%) (n=16)	1 (6.3)	13 (81.3)	2 (12.5)	0.469 <sup>a</sup>
	Incomplete, n(%) (n=16)	1 (6.3)	9 (56.3)	6 (37.5)	
	No immunization, n(%) (n=39)	2 (5.1)	26 (66.7)	11 (28.2)	
Confirmed diphtheria	Incomplete, n(%) (n=5)	1 (20.0)	3 (60.0)	1 (20.0)	0.610 <sup>b</sup>
	No immunization, n(%) (n=12)	0 (0.0)	8 (66.7)	4 (33.3)	

<sup>a</sup>Spearman's correlation coefficient (rs) = 0.087; <sup>b</sup>rs = 0.133

**Table 4.** Analysis of immunization status and fatality

Diphtheria status	Immunization status	Fatality status		P value
		Survived	Died	
Clinical diphtheria	Complete, n(%) (n=16)	15 (93.8)	1 (6.2)	0.820
	Incomplete, n(%) (n=16)	14 (87.5)	2 (12.5)	
	No immunization, n(%) (n=39)	37 (94.9)	2 (5.1)	
Confirmed diphtheria	Incomplete, n(%) (n=5)	4 (80.0)	1 (20.0)	0.294
	No immunization, n(%) (n=12)	12 (100.0)	0	

**Table 5.** Types of complications in diphtheria patients

Complication, n	(n=26)
Bullneck only	23
Bullneck, shock, epistaxis	1
Shock, epistaxis	1
Shock, myocarditis	1
Total	26

low percentage of positive culture results (17/71, 23.9%) was also noted by Puspitasari *et al.* in Dr. Soetomo General Hospital, Surabaya, Indonesia, in which only 22.8% of patients had positive culture results.<sup>14</sup> In addition, Rusmil *et al.* found that only 1.9% of subjects had positive cultures during the diphtheria outbreak in Cianjur District, West Java Province, Indonesia.<sup>15</sup> The relatively low percentage of positive culture results in

**Table 6.** Analysis of immunization status and complications

Diphtheria status	Immunization status	Complications		P value
		No complications	With complications	
Clinical diphtheria	Complete, n(%) (n=16)	14 (87.5)	2 (12.5)	0.013
	Incomplete, n(%) (n=16)	6 (37.5)	10 (62.5)	
	No immunization, n(%) (n=39)	25 (64.1)	14 (35.9)	
Confirmed diphtheria	Incomplete, n(%) (n=5)	2 (40.0)	3 (60.0)	0.620
	No immunization, n(%) (n=12)	7 (58.3)	5 (41.7)	

## Discussion

Of 71 subjects, 17 (23.9%) had positive culture results for *Corynebacterium diphtheriae*, with *mitis biovar* more common than *gravis biovar*. This finding was similar to those found in the studies of the 1990s outbreak in the newly independent states that formerly constituted the Soviet Union, in which *mitis biovar* predominated the culture results.<sup>12,13</sup> Two patients with *gravis biovar* were identified in our study. The first patient was identified in December 2011 and survived, but the second case was identified in July 2015 and died due to myocarditis. The

diphtheria studies is due to the difficulty of growing *Corynebacterium diphtheriae* bacteria in the medium.<sup>16</sup> Limited laboratory facilities also contributed to the low percentage of subjects with positive culture results.<sup>13</sup> Since the Sampang District had inadequate facilities for culturing *Corynebacterium diphtheriae*, lab cultures were done at the Surabaya Laboratory of Health (*Balai Besar Laboratorium Kesehatan Surabaya*) in Surabaya, the capital of East Java Province.

In our study, both patient groups (clinical and confirmed diphtheria) had more female than male patients. Similarly, a Hyderabad, India study

had more female than male patients, ranging from pediatric to adult patients.<sup>17</sup> A multivariate analysis by Volkze et al. in Germany found that adult females had a weaker immune response in the form of lower antibody titers which also lasted for a shorter time than those in adult males. This weaker immune response made them 45% more susceptible to diphtheria than males.<sup>18</sup>

In our study, most patients were aged 5-9 years. This finding may be due to the waning of antibody titers from inadequate booster administration.<sup>16</sup> Diphtheria also occurred mostly in early childhood years (9 years and below) due to inadequate immune system performance.<sup>16</sup> School-aged children have increased frequency of contact with the disease-causing pathogen, leading to higher natural immunity to diphtheria.<sup>19</sup>

We found no correlation between immunization status and degree of diphtheria severity. A study during the outbreak in Buri Ram, Thailand also noted the absence of such a correlation.<sup>20</sup> Various studies showed that other factors, such as residential humidity, household density, and the wall type in houses had significant correlations with the severity of diphtheria.<sup>21</sup> Sampang District is endemic to diphtheria, so residents may have frequent contact with the disease-causing pathogen. Such a situation leads to higher natural immunity to diphtheria, enabling the diphtheria severity in patients to be compromised.<sup>19</sup> We did not measure patients' antibody levels, which might have shown a direct relationship between patient immunity and diphtheria severity.

We also found no correlation between immunization status and fatality of diphtheria patients. Previous studies have shown that low immunity due to poor nutrition, younger age, inadequate or absence of immunizations, and delayed treatment may increase the risk of death in diphtheria patients.<sup>22,23</sup> However, immunization status was the sole variable, of those mentioned above (multifactorial), analyzed in this study. We also used the immunization status data from patient interviews, in addition to the Provincial Health Office data, which may have led to recall bias. Nevertheless, we consistently found that those with complete immunization status suffered less severe disease, and lower fatality, although the results were not statistically significant.

We found a statistically significant correlation between immunization status and complications in patients with clinical diphtheria, but not in those with confirmed diphtheria. The number of patients with confirmed diphtheria was significantly smaller than those with clinical diphtheria. Inadequate immunization may result in a higher risk of complications in diphtheria.<sup>20</sup> This is shown by a study of diphtheria among adult alcoholic patients in Sweden in which all patients with neurological complications had antibody titers below 0.01 IU/mL.<sup>24</sup> No similar study conducted on pediatric patients. Complete immunization was shown to increase the antibody titer in the recipients, possibly decreasing the probability of complications.<sup>19</sup> The absence of a correlation between immunization status and complications in patients with confirmed diphtheria was because none of the confirmed diphtheria cases had complete immunization status. Thus, in confirmed diphtheria cases, the risk of developing complications for incomplete and unimmunized patients was similarly higher compared to those with complete immunizations.

This study was limited as there were multiple sources of data: official medical records at the East Java Provincial Health Office, data obtained by the Diphtheria Research Team, and history-taking from diphtheria patients and their families. Data obtained from diphtheria patients and their families may have been subject to recall bias, causing inaccuracy of some parts of the data. In addition, we did not measure the antibody level against diphtheria toxin, so a more objective measurement could not be provided.

In conclusion, there is a significant correlation between immunization status and complications in patients with clinical diphtheria, but not in patients with confirmed diphtheria. This absence is likely because none of confirmed diphtheria cases have complete immunization for diphtheria.

## Conflict of Interest

None declared.

## Funding Acknowledgment

The authors received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

## References

1. Ogle JW, Anderson MS. Diphtheria in: Hay WM, Levin MJ, Deterding RR, Abzug MJ, Sondheimer JM, editor. *Current Diagnosis & Treatment Pediatrics*, 21<sup>st</sup> ed. San Francisco: McGraw-Hill; 2012. p. 1255-7.
2. Dinas Kesehatan Provinsi Jawa Timur. Profil Kesehatan Provinsi Jawa Timur Tahun 2012. Surabaya: Dinas Kesehatan Provinsi Jawa Timur; 2013. p. 34-36.
3. Dinas Komunikasi dan Informatika Jawa Timur 2014. Kasus difteri di Jatim alami penurunan. [cited 1 August 2015]. Available from: <http://kominfo.jatimprov.go.id/watch/41913>.
4. Carroll KC. *Corynebacterium diphtheriae*. In: Brooks, GF, Carroll KC, Butel JS, Morse SA, Mietzner TA, editors. *Jawetz, Melnick & Adelberg's medical microbiology*, 26<sup>th</sup> ed. San Francisco: McGraw-Hill; 2013. p. 188-190.
5. Kaufmann SHE, Sher A, Ahmed R. *Immunology of infectious disease*. Washington DC: ASM Press; 2002. p.207-221
6. Basuki PS, Soegijanto S. *Diphtheria*. In: Bagian Ilmu Kesehatan Anak RSUD Dr. Soetomo. Pedoman Diagnosis dan Terapi Bag/SMF Ilmu Kesehatan Anak Rumah Sakit Umum Dokter Soetomo Surabaya. Vol. I, ed. 3. Surabaya: Fakultas Kedokteran Universitas Airlangga; 2008. p. 76-83.
7. Cherry JD, Harrison GJ, Kaplan SL, Steinbach WJ, Hotez PJ, editors. *Felgin and Cherry's Textbook of Pediatric Infectious Diseases*. 7<sup>th</sup> ed. Philadelphia: Elsevier Saunders; 2014. p. 1301-10.
8. Acang N. *Difteri*. In: Sudoyo AW, Setiyohadi B, Alwi I, Simadibrata M, Setiati S, editor. *Buku Ajar Ilmu Penyakit Dalam*, vol. III 5<sup>th</sup> ed. Jakarta: Interna Publishing; 2010. p. 2955-61.
9. Sunarno, Sariadji K, Wibowo HA. Potensi Gen dtx dan dtxR sebagai marker untuk deteksi dan pemeriksaan toksigenitas *Corynebacterium diphtheriae*. *Buletin Penelitian Kesehatan*. 2013;41:1-10.
10. Pracoyo NE, Roselinda. Survei titer anti bodi anak sekolah usia 6-17 tahun di daerah KLB difteri dan non KLB di Indonesia. *Buletin Penelitian Kesehatan*. 2013;41:237-47.
11. Kementerian Kesehatan Republik Indonesia. Situasi Imunisasi di Indonesia. Jakarta: Pusat Data dan Informasi Kementerian Kesehatan RI; 2016. p. 1-11.
12. Dittmann S, Wharton M, Vitek C, Ciotti M, Galazka A, Guichard S, et al. Successful control of epidemic diphtheria in the states of the former Union of Soviet Socialist Republics: lesson learned. *J Infect Dis*. 2000;181:S10-22.
13. Markina SS, Maksimova NM, Vitek CR, Bogatyreva EY, Monisov AA, Diphtheria in the Russian Federation in the 1990s. *J Infect Dis*. 2000;181:S27-34.
14. Puspitasari D, Ernawati, Husada D. Gambaran klinis penderita difteri anak di RSUD Dr. Soetomo (Clinical features of children with diphtheria on Soetomo Hospital). *Media Jurnal Ners*. 2012;7:1-7.
15. Rusmil K, Chairulfatah A, Fadlyana E, Dhamayanti M. Wabah Difteri di Kecamatan Cikalang Wetan, Kabupaten Cianjur, Jawa Barat, Indonesia. *Sari Pediatri*. 2011;12(6):397-403.
16. Felgin RD, Cherry JD, editors. *Felgin and Cherry's textbook of pediatric infectious diseases*. 3<sup>rd</sup> ed. Philadelphia: Elsevier Saunders; 1992. p. 1110-1115
17. Bitragunta S, Murhekar MV, Hutin YJ, Penumur PP, Gupte MD. Persistence of diphtheria in Hyderabad, India, 2003-2006. *Emerg Infect Dis*. 2008;14:1144-6.
18. Volzke H, Kloker KM, Kramer A, Guertler L, Doren M, Baumeister SE, et al. Susceptibility to diphtheria in adults: prevalence and relationship to gender and social variables. *Clin Microbiol Infect*. 2006;12:961-7.
19. Hughes GJ, Mikhail AF, Husada D, Irawan E, Kafatos G, Bracebridge S, et al. Seroprevalence and determinants of immunity to diphtheria for children living in two districts of contrasting incidence during an outbreak in East Java, Indonesia. *Pediatr Infect Dis J*. 2015;34:1152-6.
20. Pantukosit P, Arpornsuwan M, Sookananta K. A diphtheria outbreak in Buri Ram, Thailand. *Southeast Asian J Trop Med Public Health*. 2008;39:690-6.
21. Sari PM, Zain IM. Pengaruh kondisi sanitasi rumah, status imunisasi, dan pengetahuan ibu terhadap kejadian difteri pada bayi di kota Surabaya. *J Mahasiswa Teknologi Pendidikan*. 2012;1:1-7.
22. Setiyono, Soetrisno R, Ismail D, Susatya B, Sudiantoro YE, Partatmo A, et al. Difteri pada anak: faktor-faktor yang mempengaruhi kematian. *Berita Kedokteran Masyarakat*. 1989;5:5-11.
23. Quick ML, Sutter RW, Kobaidze K, Malakmadze N, Strebel PM, Nakashidze R, et al. Epidemic diphtheria in the Republic of Georgia, 1993-1996: risk factors for fatal outcome among hospitalized patients. *J Infect Dis*. 2000;181:S130-7.
24. Bjorkholm B, Bottiger M, Christenson B, Hagberg L. Antitoxin antibody levels and the outcome of illness during an outbreak of diphtheria among alcoholics. *Scand J Infect Dis*. 1986;18:235-9.