

Effectiveness of single and double phototherapy on indirect hyperbilirubinemia in neonates

Nanda Susanti Milyana, Guslihan Dasa Tjipta, Muhammad Ali, Emil Azlin, Bugis Mardina Lubis, Pertin Sianturi, Beby Sofiani Hasibuan

Abstract

Background Hyperbilirubinemia is a common problem in full term newborns and phototherapy is the most widespread treatment for lowering bilirubin concentration in neonates. Double phototherapy could increase the effectiveness of treatment.

Objective To compare the effectiveness of single and double phototherapy and increasing spectral irradiance for decreasing serum bilirubin levels in neonates for indirect hyperbilirubinemia.

Methods An open, randomized, controlled trial was conducted at H. Adam Malik and Pirngadi Hospitals, Medan, from May to December 2009. Subjects were divided into two groups, those who received single phototherapy (n=30) and those who received double phototherapy (n=30) treatments. We included term newborns with neonatal jaundice in the first week of life. Serum bilirubin and average spectral irradiation levels were measured at baseline and after 12 hours and 24 hours of phototherapy treatment.

Results The mean total bilirubin levels of the single and double phototherapy groups at the beginning of therapy were 17.6 mg/dL (SD1.41) and 17.5 mg/dL (SD 1.32), respectively, with no significant difference between values. During the study period the sum of average spectral irradiance by double phototherapy was significantly higher than that of single phototherapy ($P < 0.05$). A significantly greater decrease in bilirubin levels was observed in the double phototherapy group at 12 hours and 24 hours of phototherapy compared to the single phototherapy group ($P = 0.001$).

Conclusion Double phototherapy is more effective than single phototherapy in reducing bilirubin levels in jaundiced newborns. [Paediatr Indones. 2011;51:316-21].

Keywords : neonatal jaundice, irradiation spectrum, phototherapy, bilirubin

Hyperbilirubinemia remains a common, important, and sometimes pathologic condition in newborns. Because of the risk of jaundice-associated neurotoxicity, management of neonatal hyperbilirubinemia is a subject of considerable discussion and debate among physicians.^{1,2} Although all infants experience some degree of hyperbilirubinemia in the first few days of life and most have some physiologic jaundice, the extent and duration varies among populations of different racial compositions or geographic distributions.^{3,4}

The jaundice in patients may be physiological or pathological. Hyperbilirubinemia is considered pathological if the time of appearance, duration, or serum bilirubin levels are determined to be significantly different from physiological jaundice.⁵ Phototherapy, although slow in effect, is now the treatment of choice by most clinicians.⁶ It uses visible light to treat hyperbilirubinemia in newborns. Phototherapy systems that simultaneously irradiate the front and

This study has been presented at PIT IV, Medan, February 22nd- 24th, 2010.

From the Department of Child Health, Medical School, University of North Sumatera, H. Adam Malik Hospital, Medan, Indonesia.

Reprint requests to: Nanda Susanti Milyana, Departement of Child Health, Medical School, University of North Sumatera, H. Adam Malik Hospital, Jl. Bunga Lau No. 17 Medan. Tel. +62-61-8361721/8365663, Fax +62-61-8361721. E-mail: nanda.susanti@yahoo.com

the back of the baby increase the dose by delivering the same amount of irradiance per square centimeter of skin to a larger skin surface area.⁷

Our study aimed to compare neonatal serum bilirubin levels and spectral irradiance for single and double phototherapy at 12 and 24 hours of treatment.

Methods

We conducted a randomized, controlled, open trial in Haji Adam Malik and Pirngadi Hospitals, Medan, Indonesia from May to December 2009. All babies admitted to the special care nursery for uncomplicated neonatal jaundice and requiring phototherapy were eligible for the study. The need for phototherapy was determined using the American Academy of Pediatrics guidelines for management of jaundice in healthy term newborns. We excluded infants with serum bilirubin levels close to the exchange transfusion limit, increased direct bilirubin, hemolytic diseases and congenital anomalies. This study was approved by the Medical Ethics Committee, University of Sumatera Utara Medical School.

Sample size needed was estimated to be 30 neonates per group. In total, 60 babies were included. Demographic and laboratory data were obtained, including age, gender, hemoglobin and albumin levels. The babies included in the study were randomized to receive either single phototherapy (control group) or double phototherapy. Subjects remained in their assigned group until after 24 hours of phototherapy. Light intensity was measured as spectral irradiance ($\mu\text{W}/\text{cm}^2/\text{nm}$) using a Dale 40 light intensity meter (USA).

Phototherapy units used were manufactured by Tessna, USA. The phototherapy units utilized five compact blue fluorescent lamps (Toshiba 20WT52),

with irradiation of $6.6 \mu\text{W}/\text{cm}^2$ by radiometer. Distance between the phototherapy unit and baby was standardized at 45 cm above and 10 cm below the baby. We used blue light photolights (Toshiba 20WT52) with irradiation of $6.6 \mu\text{W}/\text{cm}^2$ by radiometer (Dale).

Primary outcome measures were the mean differences in serum bilirubin levels at baseline, 12 hours and 24 hours. Secondary outcome measures were spectral irradiance of single phototherapy (control group) and double phototherapy. At baseline and at 12 and 24 hours of phototherapy, serum bilirubin levels and spectral irradiance were measured.

The safety of both methods was assessed and compared by monitoring body temperature, hydration status (monitored clinically and by weight measurement), skin problems (such as rashes) and gastrointestinal problems (such as loose stools or feeding intolerance).

Associations between phototherapy types and serum bilirubin levels were analyzed by Student's *t*-test. We analyzed data with SPSS version 15.0. Statistical significance was accepted as $P < 0.05$ with a 95% confidence interval.

Results

Out of 66 neonates with hyperbilirubinemia, 5 with direct hyperbilirubinemia were excluded, for a total of 61 subjects enrolled in our study (**Figure 1**). However, 1 subject dropped out due to blood sample damage.

Subjects were divided into two groups of 30 each. One group received single phototherapy and the other group received double phototherapy. Infants' characteristics, including gender, age, initial bilirubin levels before phototherapy, albumin and hemoglobin levels are shown in **Table 1**.

Table 1. Baseline characteristics

Characteristic	Double phototherapy (n= 30)	Single phototherapy (n= 30)
Gender, male/female	16/14	17/13
Mean age at phototherapy, days (SD)	5.0 (1.34)	4.9 (1.32)
Mean weight, grams (SD)	2673.3 (149.51)	2720.2 (180.83)
Mean temperature, °C (SD)	36.8 (0.35)	36.8 (0.31)
Mean initial bilirubin level, mg/dL (SD)	17.5 (1.32)	17.6 (1.41)
Mean albumin, g/dL (SD)	2.6 (0.24)	2.7 (0.25)
Hemoglobin, g/dL (SD)	14.0 (1.53)	14.0 (1.02)




Figure 1. Study profile

There were significant decreases from the initial bilirubin levels to those measured after 12 and 24 hours in the double phototherapy group. For the single phototherapy group, a significant decrease was observed from the initial to the 24 hour bilirubin level only, but not in the 12 hour measurement (Table 2).

A significantly greater decrease in bilirubin levels was observed in the double phototherapy group at 12 hours and 24 hours of phototherapy compared to the single phototherapy group ($P = 0.001$) (Table 3).

Figure 2 shows significant differences in irradiance between single and double phototherapy ($P < 0.05$) at initial, 12 hours and 24 hours of phototherapy. The total irradiation 45 cm above and 10 cm below the neonate’s body in the double phototherapy group was approximately $29.2 \mu\text{W}/\text{cm}^2/\text{nm}$.

We observed the adverse effect of hyperthermy ($T > 37.5^\circ\text{C}$) in 8 (13.3% of total) subjects from both the single and double phototherapy groups, 3 and 5 subjects, respectively. Other potential side effects, such as diarrhea and dehydration, were not found during monitoring.

Table 2. Mean decreases in serum total bilirubin levels between initial and 12 hours, and between initial and 24 hours in the double and single phototherapy groups

Type of phototherapy	Mean (SD)	95% CI	P
Double phototherapy			
Decrease in bilirubin from baseline to 12 hours (mg/dL)	6.5 (0.63)	6.3 to 6.7	0.001
Decrease in bilirubin from baseline to 24 hours (mg/dL)	10.1 (1.05)	9.7 to 10.4	0.001
Single phototherapy			
Decrease in bilirubin from baseline to 12 hours (mg/dL)	0.1 (0.25)	0.0 to 0.1	0.059
Decrease in bilirubin from baseline to 24 hours (mg/dL)	3.8 (1.32)	3.5 to 4.5	0.001

Table 3. Bilirubin levels at baseline, and after 12 and 24 hours of phototherapy

	Double phototherapy Mean (SD)	Single phototherapy Mean (SD)	95% CI	P
Initial bilirubin	17.5 (1.32)	17.6 (1.41)	- 5.8 to 0.8	0.694
Bilirubin 12 hours	11.0 (1.43)	17.6 (1.44)	5.8 to 7.3	0.001
Bilirubin 24 hours	7.4 (1.61)	13.8 (1.84)	7.4 to 9.0	0.001




Figure 2. Comparison of spectral irradiance in single and double phototherapy at initial, 12 hours and 24 hours of phototherapy

Discussion

The average age of subjects in our study was 4 to 5 days. It is known that elevated bilirubin levels often peak in the first week of life.³ Hyperbilirubinemia is a serious, and potentially life-threatening condition. This condition is the main reason for hospital revisits by full term infants in their first week of life.⁸

A significant decrease of bilirubin level between baseline and 12 hours and baseline and 24 hours was seen in the double phototherapy group, with an average decrement of 6.5 mg/dl ($P=0.001$) and 10.1 mg/dL ($P=0.001$), respectively. For the single phototherapy group, a significant decrease in bilirubin level was observed only between baseline and 24 hours, with an average decrement of 3.8 mg/dL ($P= 0.001$).

Double phototherapy using blue light with wavelength 430-490 nm and spectral irradiance of $\geq 30 \mu\text{W}/\text{cm}^2/\text{nm}$ (measured with radiometer or estimated by placing the baby directly under the light or widening the exposed surface) is effective for reducing bilirubin levels.^{9,10} Double phototherapy was more effective than single phototherapy in our study. Newborns receiving double phototherapy had a larger surface area exposed to constant irradiance thereby increasing their total dose of phototherapy compared to those undergoing single phototherapy. This increased dose causes the production of more lumirubin.¹¹ However, merely increasing the surface

area exposed, while maintaining the same total energy output to the skin, does not improve the efficacy of phototherapy.^{12,13} A randomized, clinical trial in Thailand showed that double phototherapy was safer and more effective in reducing bilirubin levels compared to single phototherapy.¹⁴ Double phototherapy is an alternative mode of intensive phototherapy that is effective, economical and easy to use.

A study in Brazil compared the effectiveness of double phototherapy with total irradiance $75.6 \mu\text{W}/\text{cm}^2/\text{nm}$ and pharmacotherapy, and found that double phototherapy was better and safer in reducing bilirubin levels with minimal side effects.¹⁵ Similar studies, including one in Saudi Arabia, also showed that double phototherapy with blue light was more effective than single phototherapy, with minimal side effects.¹⁶⁻¹⁸

Light intensity is the factor that determines the effectiveness of phototherapy. Higher light intensity reduced bilirubin levels faster. A study in England showed that phototherapy with maximum irradiance and wide exposure shortened the duration of phototherapy.²²

In our study, the nearest light source was 10 cm from the neonates' bodies. During phototherapy, the eyes and genitalia were covered to avoid damage from exposure to high light intensity.¹⁹⁻²¹

Blue light of wavelength 425-475 nm is the best light type to reduce indirect bilirubin levels.^{23,24} Blue light is very effective because it has a shorter

wavelength compared to other visible light, with the exception of purple. Wavelength is inversely proportional to energy level; the shorter the wavelength, the greater energy produced.²⁴

In our study we used blue light. Light intensity was measured at baseline, and at 12 and 24 hours of phototherapy. There was a significant difference in spectral irradiance between the two groups at all time points ($P = 0.001$). To increase photolight intensity for both groups, we placed the neonates supine and varied positions every 3 hours during phototherapy. However, a clinical trial in Israel comparing 14 neonates in alternating positions to 16 others in a supine position while using single phototherapy showed a significantly greater decrease in bilirubin levels in the supine group after 24 hours of phototherapy.²⁵

Neonates treated for high bilirubin levels can also suffer from dehydration and may require additional fluid intake.²⁶ Neonatal maturity, adequate caloric intake, photolight unit temperature, distance between the neonate and photolight, and incubator rate of heat loss are all potential factors in increasing neonatal body temperature, environmental temperature, insensible water loss, as well as respiratory rate and blood flow to the skin. Increased peripheral blood flow can increase fluid loss, requiring adjustment by administration of intravenous fluids.^{26,27} Changes in skin, such as rash, darker skin colour and burning can be seen if infants are overexposed to fluorescent light. A study in the Netherlands found that during intensive phototherapy, a 20% increment of total fluid requirement may prevent increased body temperature.²⁸

Body temperature and fluid administration were strictly monitored. Fluid intake was given every 2 hours, and was increased by 10 - 20% of the total fluid requirement. In breastfed neonates, phototherapy was withheld during breastfeeding.^{29,30} In our study, we found hyperthermia ($T > 37.5^{\circ}\text{C}$) in 3 (5% of total) neonates in the single phototherapy group and 5 (8.3% of total) in the double phototherapy group.

A limitation of our study was not including maternal characteristic data associated with hyperbilirubinemia in neonates.

In conclusion, we found a significantly greater decrease in bilirubin levels in the double phototherapy group at 12 hours and 24 hours of

phototherapy compared to the single phototherapy group. Double phototherapy is more effective than single phototherapy in reducing bilirubin levels in jaundiced newborns.

References

1. Sukadi A. Hiperbilirubinemia. In: Kosim MS, Yunanto A, Dewi R, Sarosa GI, Usman A, editors. Buku ajar neonatologi. 1st ed. Jakarta: IDAI; 2008. p.147-69.
2. Martiza L. Ikterus. In: Juffrie M, Oswari H, Arief S, Rosalina I, editors. Buku ajar gastroenterologi-hepatologi. 1st ed. Jakarta. Badan Penerbit IDAI; 2010. p.263 – 84.
3. Martin CR, Cloherty JP. Neonatal hyperbilirubinemia. In: Cloherty JP, Eichenwald EC, Stark AR, editors. Manual of neonatal care. 5th ed. Philadelphia: Lippincott, Williams & Wilkins. 2004; p. 185-219.
4. Dennery AP, Seidman DM, Stevenson KD. Neonatal hyperbilirubinemia. *N Engl J Med*. 2001;8:581-90.
5. Hammerman C, Kaplan M. Recent developments in the management of neonatal hyperbilirubinemia. *NeoReviews*. 2000;1:19.
6. Stoll BJ, Kliegman RM. Jaundice and hyperbilirubinemia in the newborn. In: Behrman RE, Kliegman RM, Jenson HB, editors. Nelson textbook of pediatrics. 17th ed. Philadelphia: Elsevier Saunders; 2006. p. 592-98.
7. Madan A, Macmahon JR, Stevenson DK. Neonatal hyperbilirubinemia. In: Taeusch HW, Ballard RA, Gleason CA, editors. Avery's disease of the newborn. 8th ed. Philadelphia: Elsevier Saunders; 2005. p. 1226-53.
8. Sarici SU, Alpay F, Unay B, Ozcan O, Gokcay E. Double versus single phototherapy in term newborn with significant hyperbilirubinemia. *J Trop Pediatrics*. 2000;46:36-9.
9. Holtrop PC, Ruedisueli K, Maisels MJ. Double versus single phototherapy in low birth weight newborns. *Pediatrics*. 2008;90:674-7.
10. Tan KL. Efficacy of bidirectional fiber-optic phototherapy for neonatal hyperbilirubinemia. *Pediatrics*. 1997;99:e13.
11. Erika R, Harianto A, Indarso F, Damanik MS. Hiperbilirubinemia pada neonatus. 2007 Dec [cited 2008 November]. Available from: <http://www.pediatrik.com/pkb/20060220-js9>.
12. Sarici SU, Serdar MA, Korkmaz A, Erdem G, Oran O, Tekinalp G, et al. Incidence, course and prediction of hyperbilirubinemia in near-term and term newborn. *Pediatrics*. 2004; 113:775-80.
13. Kaplan M, Muraca M, Hammerman C, Rubaltelli FF, Vilei MT, Vreman HJ, et al. Imbalance between production

- and conjugation of bilirubin: a fundamental concept in the mechanism of neonatal jaundice. *Pediatrics*. 2002;13:110.
14. Boonyarittipong P, Kriangburapa W, Booranavanich K. Effectiveness of double-surface intensive phototherapy versus single-surface intensive phototherapy for neonatal hyperbilirubinemia. *J Assoc Thai*. 2008;90:50-5.
 15. Facchini FP, Bianchi MO, Silva BA. Intensive phototherapy treatment for severe haemolytic disease of the newborn. *J Pediatr (Rio J)*. 2000;76:387-90.
 16. Al-Alaiyan S. Fiberoptic, conventional and combination phototherapy for treatment of non-hemolytic hyperbilirubinemia in neonates. *Ann Saud Med*. 1996;16:633-6.
 17. Sarin M, Dutta S, Narang A. Randomized controlled trial of compact fluorescent lamp versus standard phototherapy for the treatment of neonatal hyperbilirubinemia. *Indian Pediatr*. 2006;43:583-90.
 18. Garg AK, Prasad RS, Al-Hifzi I. A controlled trial of high-intensity double-surface phototherapy on fluid bed versus conventional phototherapy in neonatal jaundice. *Pediatrics*. 1995;95:914-16.
 19. Knudsen A, Brodersen R. Skin colour and bilirubin in neonates. *Arch Dis Child*. 1998;64:605-9.
 20. Subcommittee on hyperbilirubinemia. American Academy of Pediatrics. Management of hyperbilirubinemia in the newborn infant 35 or more weeks of gestation. *Pediatrics*. 2004;114:297-316.
 21. Maisels MJ, Donagh FA. Phototherapy for neonatal jaundice. *N Engl J M*. 2008;358:920-8.
 22. Eggert P, Stick C, Schroder H. On the distribution of irradiation intensity in phototherapy. Measurements of effective irradiance in an incubator. *Eur J Pediatr*. 1984;142:58-61.
 23. Hobbie R, Roth B. Atoms and light. 2007 Oct 13 [cited 2008 November 10]. Available from: <http://www.springerlink.com/viewarticle>.
 24. Stokowski LA, Short MRN, Witt CL. Fundamentals of phototherapy for neonatal jaundice. 2006 Dec [cited 2008 November 10]. Available from: <http://www.emedicine.com/viewarticle/551363>.
 25. Shinwell ES, Sciaky Y, Karplus M. Effect of position changing on bilirubin levels during phototherapy. *J of Perinatol*. 2002;22:226-9.
 26. Gomella TL. Hyperbilirubinemia indirect. In: Gomella TL, editor. *Neonatology: management, procedures, on-call problems, disease, and drugs*. 5th ed. New York: McGraw-Hill; 2004. p. 247-50.
 27. Kang JH, Shankaran S. Double phototherapy with high irradiance compared with single phototherapy in neonates with hyperbilirubinemia. *Am J Perinat*. 1995;12:178-80.
 28. Grunhagen DJ, De Boer MGJ, Beaufort AJ. Transepidermal water loss during halogen spotlight phototherapy in preterm infants. *Pediatr Res*. 2002;51:402-5.
 29. Bagchi A. Phototherapy. In: MacDonald MG, Ramasethu J, editors. *Procedures in neonatology*. 3rd ed. Philadelphia: Lippincott William Wilkins; 2002. p. 373-8.
 30. Hansen TWR. Phototherapy for neonatal jaundice-therapy effects on more than one level?. *Sem in Perinatol*. 2010;34:231-34.